




supporo COMPRESSION

MEDICAL SIZE - CHOOSING THE CORRECT SIZE:



- Obtain patient's measurements from a trained health care professional
- Measurements should be taken early in the day before swelling occurs
- Measurements should not be done by the patient
- Determine style preference (length: knee or thigh high; open or closed toe, colour)
- Determine height measurements to choose whether require regular or long type. Measure A to D for knee high; A to F or G for thigh high.
- Determine circumference measurements at points B, D, and F depending on the style desired;

A = floor
B = ankle

D = mid calf at largest circumference
F = mid point between knee and G
G = groin area

Consult chart with emphasis on ankle (B) position if other measurement points (D, F, G) suggest a different size

Long or short length

LENGTH / LONGUEUR

	SHORT / COURT		LONG	
	inches / pouces	cm	inches / pouces	cm
A-D	13 - 15	34 - 38	15 - 17	39 - 44
A-F	20 - 23	52 - 59	23 - 27	60 - 68
A-G	24 - 28	62 - 71	28 - 32	72 - 82

Circumference measurements

CIRCUMFERENCE / CIRCONFÉRENCE

	SMALL / PETIT					MEDIUM					LARGE / GRAND					X-LARGE / T-GRAND					XX-LARGE / TT-GRAND			
	inches / pouces		cm			inches / pouces		cm			inches / pouces		cm			inches / pouces		cm			inches / pouces		cm	
	normal	max	normal	max		normal	max	normal	max		normal	max	normal	max		normal	max	normal	max		normal	max	normal	max
G	16-18	18-20	42-46	47-51	G	18-20	20-22	47-51	52-57	G	20-22	23-25	52-57	58-63	G	23-25	25-28	58-63	64-70	G	25-28	28-31	64-70	71-78
F	14-16	16-18	37-40	41-45	F	16-18	18-20	41-45	46-50	F	18-20	20-22	46-50	51-56	F	20-22	22-25	51-56	57-62	F	22-25	25-27	57-62	63-69
D	11-12	12-13	28-30	31-34	D	12-13	14-15	31-34	35-38	D	14-15	16-17	35-38	39-42	D	16-17	17-19	39-42	43-47	D	17-19	19-21	43-47	48-52
B	6,6 - 8,2	17 - 21	B	8,2 - 9,8	21 - 25	B	9,8-11,4	25 - 29	B	11,4-12,9	29 - 33	B	12,9-14,5	33 - 37										